

Exercises After Foot or Toe Injury

Notes:

This pamphlet is for educational purposes only. It is not intended to replace the advice or professional judgment of a health care provider. The information may not apply to all situations. If you have any questions, please ask your health care provider.

Find this pamphlet and all our patient resources here:
<https://library.nshealth.ca/Patients-Guides>

Connect with a registered nurse in Nova Scotia any time:
Call 811 or visit: <https://811.novascotia.ca>

Prepared by: Physiotherapy
Designed by: Nova Scotia Health Library Services

WB85-1451 © December 2024 Nova Scotia Health Authority
To be reviewed December 2027 or sooner, if needed.

Exercises After Foot or Toe Injury

You may now move your foot and/or toe. You may feel discomfort when you first start the exercises. This is common.

You may have some bruising or swelling in your ankle and/or foot. This will get better over time.

What will happen after my cast/bandage is taken off?

Swelling

- Your foot may be bruised or swollen.
- Swelling may last for months.
- The amount of swelling you have may change throughout the day.

To lower swelling:

- Raise your leg when you are resting, so that your foot is above the level of your heart.
- Bend your ankle and point your foot often.
- If you have been given a compression stocking:
 - › Wear it during the day.
 - › Take it off to do your exercises and to sleep.

Balance exercises

□ Exercise 10

- Stand facing something stable (like a kitchen counter or chair) and hold on with both hands.
- Stand on your affected foot. Hold for 15 to 20 seconds. Relax.
- Repeat _____ times.



Progressions

- As your leg gets stronger, lower the amount of support you use to help you balance on 1 leg.

Try:


- › Fingertip support with 2 hands
- › Using only 1 hand for support
- › Not using any hands for support
- When your balance is better and you can balance on 1 leg with no support, try balancing with your eyes closed.

□ Exercise 9

- Stand facing stable something stable (like a kitchen counter or chair) and hold on with both hands.
- Use both of your feet to rise up onto your toes. Hold for 5 seconds, then slowly lower.
- Repeat _____ times.

Progression:

- When you are feeling stronger, stand only on your affected foot and rise up on your toes.
- Hold for 5 seconds, then slowly lower.
- Repeat _____ times.



- Raise your foot at night by putting pillows under your leg or by putting a pillow under your mattress at the end of the bed.
- Use ice (see page 3).
- Start massaging your scar when told by your physiotherapist. Your incision (cut) scar must be well-healed.

Stiffness

- Your foot and/or toe may be stiff (hard to move). This is common.
- To help get the movement back in your joints, start moving your foot and/or toe as told by your physiotherapist.
- You may feel discomfort when you first start the exercises. This will go away over time, as your foot and/or toes become more flexible.

Walking

- Your doctor or physiotherapist will tell you:
 - › how much weight you may put on your foot.
 - › the type of walking aid to use (like a cane or crutches), if needed.
- Try to walk a bit more each day.

Heat and ice

- Before exercising, use heat (like a hot pack, a hot water bottle, or a soak in warm water) for about 15 minutes to help loosen up your foot and/or toe.
 - › Your incision **must** be well-healed and not infected before using heat or soaking in water.
- Use ice (like a gel pack or a bag of frozen vegetables) for about 15 minutes after exercising to help control pain in your foot and/or toe.

Spacer

If you have a toe spacer, take it off to do your exercises.

Exercises

- It is important to hold the stretches for the full amount of time. This helps your muscles stretch.
- **Only do the exercises you have been asked to do.** See which exercises (starting on next page) have been checked off by your physiotherapist.
- **Plan to do your exercises 3 times each day.**

Strengthening exercises

Exercise 7

- Put a towel on the floor in front of a chair. Sit in the chair.
- Place your heel on the floor and the ball of your affected foot on the towel.
- Grasp the towel with your toes, bunching the towel. Hold for 5 seconds. Relax.
- Repeat _____ times.



Exercise 8

- Sit in a chair.
- Raise your heel up while keeping your toes flat on the floor, and pushing your toes down. Hold for 5 seconds.
- Slowly lower your heel back down and relax.
- Repeat _____ times.



☐ Exercise 6

- Sit in a chair.
- Bend the knee of your affected leg so that you can reach your toes. Hold the end of your toe.
- Slowly bend the **tip** of your toe downwards until you feel a stretch. Hold for 10 to 30 seconds. Relax.
- Repeat _____ times.



Exercises

☐ Exercise 1

- Move your toes up. Hold for 5 seconds. Relax.
- Repeat _____ times.



☐ Exercise 2

- Bend your toes down. Hold for 5 seconds. Relax.
- Repeat _____ times.



What are your questions?

Please ask. We are here to help you.

□ Exercise 3

- Move your big toe out to the side, away from the rest of your toes. Hold for 5 seconds. Relax and move your big toe back to the starting position.
- Repeat _____ times.



Stretching exercises

□ Exercise 4

- Sit in a chair.
- Bend the knee of your affected leg so that you can reach your toes.
- Keep the bottom of your toes still with 1 hand. Use your other hand to hold the end of your toe.
- Slowly bend your toe downwards until you feel a stretch. Hold for 10 to 30 seconds. Relax.
- Repeat _____ times.



□ Exercise 5

- Sit in a chair.
- Bend the knee of your affected leg so that you can reach your toes. Hold the end of your toe.
- Slowly bend your toe upwards until you feel a stretch. Hold for 10 to 30 seconds. Relax.
- Repeat _____ times.

